



HHS Public Access

Author manuscript

JAMA. Author manuscript; available in PMC 2016 September 15.

Published in final edited form as:

JAMA. 2016 March 15; 315(11): 1149–1158. doi:10.1001/jama.2016.1353.

Association Between Vaccine Refusal and Vaccine-Preventable Diseases in the United States:

A Review of Measles and Pertussis

Varun K. Phadke, MD, Robert A. Bednarczyk, MS, PhD, Daniel A. Salmon, MPH, PhD, and Saad B. Omer, MBBS, MPH, PhD

Division of Infectious Diseases, Emory University, Atlanta, Georgia (Phadke); Department of Epidemiology, Emory University, Atlanta, Georgia (Bednarczyk, Omer); Emory Vaccine Center, Emory University, Atlanta, Georgia (Bednarczyk, Omer); Institute for Vaccine Safety, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland (Salmon); Department of Pediatrics, School of Medicine, Emory University, Atlanta, Georgia (Omer); Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, Georgia (Omer)

Abstract

IMPORTANCE—Parents hesitant to vaccinate their children may delay routine immunizations or seek exemptions from state vaccine mandates. Recent outbreaks of vaccine-preventable diseases in the United States have drawn attention to this phenomenon. Improved understanding of the association between vaccine refusal and the epidemiology of these diseases is needed.

OBJECTIVE—To review the published literature to evaluate the association between vaccine delay, refusal, or exemption and the epidemiology of measles and pertussis, 2 vaccine-preventable diseases with recent US outbreaks.

EVIDENCE REVIEW—Search of PubMed through November 30, 2015, for reports of US measles outbreaks that have occurred since measles was declared eliminated in the United States (after January 1, 2000), endemic and epidemic pertussis since the lowest point in US pertussis

Corresponding Author: Saad B. Omer, MBBS, MPH, PhD, Rollins School of Public Health, Emory University, 1518 Clifton Rd NE, Room 7017 (CNR Bldg), Atlanta, GA 30322 (somer@emory.edu).

Role of the Funder/Sponsor: The sponsor had no role in the collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Disclaimer: The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Allergy and Infectious Diseases or the National Institutes of Health.

Submissions: We encourage authors to submit papers for consideration as a Review. Please contact Edward Livingston, MD, at Edward.livingston@jamanetwork.org or Mary McGrae McDermott, MD, at mdm608@northwestern.edu.

Author Contributions: Dr Phadke had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Salmon, Omer.

Acquisition, analysis, or interpretation of data: Phadke, Bednarczyk, Omer.

Drafting of the manuscript: Phadke, Bednarczyk, Salmon, Omer.

Critical revision of the manuscript for important intellectual content: Phadke, Bednarczyk, Omer.

Statistical analysis: Phadke, Bednarczyk, Omer.

Administrative, technical, or material support: Salmon, Omer.

Study supervision: Omer.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Dr Salmon reports receiving grant funding from Crucell, Pfizer, and Merck and personal fees from Parents of Kids with Infectious Diseases. No other disclosures are reported.

incidence (after January 1, 1977), and for studies that assessed disease risk in the context of vaccine delay or exemption.

FINDINGS—We identified 18 published measles studies (9 annual summaries and 9 outbreak reports), which described 1416 measles cases (individual age range, 2 weeks–84 years; 178 cases younger than 12 months) and more than half (56.8%) had no history of measles vaccination. Of the 970 measles cases with detailed vaccination data, 574 cases were unvaccinated despite being vaccine eligible and 405 (70.6%) of these had nonmedical exemptions (eg, exemptions for religious or philosophical reasons, as opposed to medical contraindications; 41.8% of total). Among 32 reports of pertussis outbreaks, which included 10 609 individuals for whom vaccination status was reported (age range, 10 days–87 years), the 5 largest statewide epidemics had substantial proportions (range, 24%–45%) of unvaccinated or undervaccinated individuals. However, several pertussis outbreaks also occurred in highly vaccinated populations, indicating waning immunity. Nine reports (describing 12 outbreaks) provided detailed vaccination data on unimmunized cases; among 8 of these outbreaks from 59% through 93% of unvaccinated individuals were intentionally unvaccinated.

CONCLUSIONS AND RELEVANCE—A substantial proportion of the US measles cases in the era after elimination were intentionally unvaccinated. The phenomenon of vaccine refusal was associated with an increased risk for measles among people who refuse vaccines and among fully vaccinated individuals. Although pertussis resurgence has been attributed to waning immunity and other factors, vaccine refusal was still associated with an increased risk for pertussis in some populations.

Vaccine coverage, estimates of the number of people who have received vaccines, remains high among young children for most routine vaccines. For example, in 2013 national coverage was 91.9% for 1 dose or more of measles, mumps, and rubella vaccine, and 83.1% for 4 doses or more of diphtheria, tetanus, and acellular pertussis (DTaP) vaccine.¹ Many childhood vaccine-preventable diseases have been effectively controlled.² However, recent outbreaks of vaccine-preventable diseases in the United States have prompted clinicians, public health officials, politicians, the media, and the public to pay greater attention to the growing phenomenon of vaccine refusal and hesitancy.^{3–5}

In individual studies, vaccine refusal has been associated with outbreaks of invasive *Haemophilus influenzae* type b disease,⁶ varicella,⁷ pneumococcal disease,⁸ measles,⁹ and pertussis.¹⁰ The most direct measure of vaccine refusal is the rate of parents claiming nonmedical exemptions to school immunization requirements. Nonmedical exemptions can include religious exemptions, if a parent feels that immunizations conflict with their religious or spiritual beliefs (eg, objection to the use of fetal tissue in the production of some vaccines), or personal belief exemptions, if a parent objects to immunizations for moral or philosophical reasons (eg, objection to the use of nonnatural products or the total number of vaccines to be administered). Sometimes it is difficult to distinguish between purely religious or philosophical objections to immunization and safety concerns about vaccines that manifest as nonmedical exemptions.^{3–5} Over the past 20 years, rates of nonmedical exemptions have steadily increased.^{10,11} Initially this increase in exemptions was seen only among states that more readily offered personal belief exemptions (ie, those with easier procedures for obtaining such exemptions).¹⁰ However, more recently, states with

moderately difficult exemption procedures and religious exemptions have experienced increases in rates.¹²

An outbreak of measles in late 2014 that originated at Disneyland in Anaheim, California, highlighted vaccine refusal and related disease outbreaks.¹³ This outbreak was associated with 111 cases (accounting for two-thirds of the total measles cases reported by April 2, 2015) in 7 US states, Canada, and Mexico.^{14,15} Approximately half the cases were among unvaccinated persons, most of whom were eligible for vaccination yet intentionally remained unvaccinated.^{14,16} Although previous studies demonstrated associations between rates of nonmedical exemption and the risk of acquiring vaccine-preventable diseases, they have been of varying design (eg, modeling studies) and quality, and did not systematically examine measles (or other vaccine-preventable diseases) and the contribution of refusals to disease outbreaks. This review evaluates the available evidence to characterize the relationship between vaccine refusal and the epidemiology of measles and pertussis, 2 vaccine-preventable diseases with recent outbreaks in the United States.

Methods

Measles

We first searched for US-based measles outbreaks using PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/>) for studies published through November 30, 2015, using the following search terms: *measles vaccinated "United States," measles unvaccinated "United States," measles outbreak "United States," measles import* "United States," measles case* "United States,"* and *measles MMWR*. We restricted our search to after the measles elimination era (reports published on or after January 1, 2000¹⁷) to provide the most relevant review of recent measles epidemiology. To evaluate the risk of measles associated with exemptions to vaccine mandates, we also searched using the terms: *measles risk exemption*. For this latter search we used no date restriction because we wanted to capture studies that directly assessed risk related to vaccine exemption, regardless of whether the outbreak occurred before or after elimination.

Pertussis

Unlike measles, pertussis remains endemic in the United States, and its incidence has been steadily rising since it reached a nadir in 1976.¹⁸ To account for this important difference between measles and pertussis, we adapted the search strategy for pertussis as follows: we first searched for US-based pertussis outbreaks, this time restricting our search to reports of nonoverlapping community or state pertussis outbreaks that have occurred since the lowest point in US pertussis incidence (those published after January 1, 1977). We again searched Pub Med for studies published through November 30, 2015, using the following search terms: *pertussis epidemic, pertussis endemic, pertussis outbreak, and pertussis MMWR*. To evaluate the risk of pertussis associated with vaccine exemption, we also searched using the terms: *pertussis vaccine refusal, pertussis vaccine exemption, pertussis vaccine delay, pertussis vaccine hesitan*, pertussis undervaccination, and pertussis cluster*. For this latter search we again used no date restrictions to capture studies that may have been published

earlier than the start of the pertussis resurgence, because we expected any association between vaccine refusal and disease risk to be similar.

Results

Measles

Recent Epidemiology of Measles in the United States—Measles-containing vaccines were first licensed in the United States in 1963. Following their widespread introduction and 2 intensive elimination efforts over the following 2 decades, the incidence of measles fell dramatically, from more than 300 cases per 100 000 population in the prevaccine era to a median 1.3 cases per 100 000 from 1982 through 1988.¹⁹ From 1989 through 1992 there was a resurgence of measles in multiple countries, including the United States, thought to be primarily due to suboptimal vaccine coverage among preschool-aged children, as well as a small number of vaccine failures after a single dose of measles-containing vaccine. This motivated an intense public health effort to improve vaccine coverage in this age group.²⁰ A third elimination effort, which started in the 1990s and included formal introduction of a second dose of measles-containing vaccine, further reduced the incidence of measles to less than 1 case per 1 million population by 1997–1999, culminating in successful elimination of endemic measles in the United States by 2000.¹⁷ Since then, there have been a number of measles outbreaks in the United States, without reemergence of indigenous transmission. These importation-related outbreaks (eg, linked to individuals who acquired measles outside the United States and introduced it into their community upon return) totaled fewer than 250 measles cases annually until 2014, when the United States experienced 23 measles outbreaks and the Centers for Disease Control and Prevention (CDC) reported 668 measles cases, the highest number recorded since elimination (Box 1).¹⁵

Box 1

Recent Epidemiology of Measles and Pertussis in the United States

Measles

- 1963: Measles-containing vaccines are first licensed in the United States
- 1988: Following the introduction of measles-containing vaccines and 2 measles elimination efforts, measles incidence falls to 1.3 cases per million population
- 1989–1992: Measles resurgence in multiple countries, including the United States, attributed to suboptimal vaccine coverage in preschool-aged children
- 1989: Introduction of a routine second dose of measles-containing vaccines for preschool-aged children and beginning of third elimination effort
- 2000: Measles is declared eliminated from the United States

- 2014: 23 measles outbreaks—including 1 that originated in Disneyland in Anaheim, California—and 668 measles cases are reported in the United States, the largest annual total in the era after elimination

Pertussis

- 1940s: Whole-cell pertussis vaccines come into widespread use in the United States
- 1976: Pertussis incidence reaches a nadir, with only 1010 cases reported in the United States
- 1990s: Acellular pertussis vaccines replace whole-cell vaccines in the immunization schedule
- 2004: 25 827 cases of pertussis are reported in the United States, the largest number since 1959
- 2005: Acellular pertussis vaccines are licensed for use in adults and adolescents (tetanus, reduced diphtheria, and acellular pertussis [Tdap] vaccine)
- 2010: Epidemic of pertussis in California that included 9154 reported cases, with the highest incidence of disease in infants younger than 6 months, followed by children aged 7–10 years; in the latter age group, disease risk was associated with an increasing interval since the last dose of pertussis-containing vaccine
- 2012: Tdap vaccine administration is recommended during every pregnancy
- 2014: Epidemic of pertussis in California that included 9935 reported cases

Vaccine Refusal and Measles Outbreaks—Using the search strategy for measles outbreaks, a total of 687 articles were identified (eFigure 1A in the Supplement), of which 65 were reports of measles outbreaks between 2000 and 2015. Of these 65 reports, we excluded 34 that were either duplicate reports (eg, a *JAMA* republication of an *Morbidity and Mortality Weekly Report [MMWR]* article) or that contained data that were updated or summarized in later reports. Of the remaining 31 reports, we selected 18 to generate a summary of measles outbreaks in the United States from 2000 through 2015 (Table 1).^{14,21–37} Nine of these reports are surveillance summary reports from the CDC, summarizing multiple outbreaks during the reporting period.^{14,21–24,27,33,34,36} The remaining 9 reports are investigations related to a specific measles outbreak or importation.^{25,26,28–32,35,37} The 13 reports not used in the summary of measles outbreaks described individual outbreaks whose case data were captured and/or updated in the other 18 summary reports.

Since January 1, 2000, there were 1416 reported measles cases (individual age range, 2 weeks–84 years; 178 cases younger than 12 months) in the United States associated with

outbreaks or periodic summary reports published through November 30, 2015.^{14,21–37} A total of 199 cases (14.1%) involved individuals with a history of receiving a measles-containing vaccine, whereas 804 cases (56.8%) involved individuals with no history of receiving a measles-containing vaccine. Additionally, there were 231 cases (16.3%) that involved individuals with unknown history of measles-containing vaccine receipt and 174 cases (16.3%) that involved individuals for whom the reported measles vaccination status did not distinguish between unknown or unvaccinated (eg, reported as “unknown or unvaccinated”). In 7 of these outbreak reports and annual summaries,^{14,29,33,34,36–38} there was sufficient information to determine the number of unvaccinated individuals with nonmedical exemptions. In these 7 reports, there were 970 cases, of which 574 cases occurred in unvaccinated individuals who were age-eligible for measles vaccination, and 405 of the unvaccinated individuals (70.6%) were unvaccinated due to nonmedical exemptions (comprising 41.8% of the total cases reported).

There were 18 outbreaks that had sufficient data (ie, day of symptom onset and vaccination status for measles cases) to construct a cumulative epidemic curve (Figure 1).^{23,26,29,31,35,37–49} These outbreaks consisted of 145 cases (median cases per outbreak, 6.5 [range, 1–22]); of these cases, 111 individuals were unvaccinated, 11 individuals had unknown vaccination status, and 23 individuals had received at least 1 dose of vaccine. These 145 cases included index cases (ie, the first case to be identified in an outbreak) as well as additional cases that were identified in the course of each individual outbreak investigation. Each outbreak investigation classified individual cases within a specific “generation of spread” based on an established chain of transmission, or by inferring the sequence of transmission based on the date of symptom onset. For example, individuals that developed measles following exposure to the index case were considered to be in the first generation of outbreak spread, those who developed measles following exposure to those in the first generation were considered to be in the second generation of spread, and so on. The outbreaks evaluated in the cumulative epidemic curve included cases that occurred up to 5 generations of spread after the index case, with the latest related case occurring 12 weeks after identification of the index case. When viewed by week of outbreak, unvaccinated individuals constituted a larger fraction of the total measles cases per week in the earliest weeks of an outbreak (eg, earlier generations).

Vaccine Refusal and Measles Risk—We identified 12 articles using the search for studies evaluating measles risk in the context of vaccine exemption (eFigure 1B in the Supplement). Of these, 10 were excluded: 1 was not published in English, 1 was an outbreak investigation, 5 were vaccine coverage analyses, 1 was a surveillance summary, and 2 were review articles related to the topic. Two studies directly examined the individual and community risks of measles associated with intentionally unvaccinated children (Table 2).^{9,50} Both were retrospective cohort studies conducted using measles case data from the era before measles elimination (before 2000), and were therefore not included in our analysis of recent measles epidemiology. The first study used national measles surveillance data reported to the CDC from 1985 through 1992, inclusive of the 1989–1992 measles resurgence.⁹ In this study, unvaccinated children who had a vaccine exemption were 35 times more likely to contract measles compared with vaccinated children. The epidemic

curve of measles determined that the resurgence started a year earlier among children with exemptions compared with vaccinated children.⁹ The investigators also used mathematical modeling to determine that the prevalence of exemptions in a geographic region was also associated with disease risk in the nonexempt population in that region, with high local aggregation of individuals with exemptions (ie, clustering) being associated with greater measles incidence. A second study used Colorado measles cases from 1987 through 1998 and found that children with exemptions were 22 times more likely to contract measles than vaccinated children.⁵⁰ The absolute and relative risk of disease was highest among individuals with exemptions aged 3 to 10 years. This study also found that the frequency of exemptions at the county level was associated with county measles incidence, though there was no association between schools with higher rates of exemptions and school-based measles outbreaks. Exemptions were more likely to be in the index case and first generation (eg, individuals that acquired measles following exposure to the index case) (14.5%) compared with later generations (eg, those who acquired measles from individuals in the first generation or later rather than the index case) (7.1%). Among vaccinated children who contracted measles, 11% contracted it from an individual with an exemption (Box 2).⁵⁰

Box 2

Vaccine Refusal and the Epidemiology of Measles and Pertussis in the United States

Measles

- Since measles was declared eliminated from the United States (eg, 2000), 1416 measles cases have been reported in published summaries or outbreak reports, and more than half (56.8%) of these cases had no history of measles vaccination
- Among the unvaccinated, age-eligible measles cases for whom the reason for nonvaccination was available, 405 (70.6%) had a nonmedical exemption to vaccination
- Unvaccinated individuals made up a greater proportion of measles cases in the index or first generation of a cumulative epidemic curve of 18 measles outbreaks
- Children with vaccine exemptions have a substantially greater risk for acquiring measles than fully vaccinated children—in 1 study the risk was 35 times that of the vaccinated population
- Higher rates of vaccine exemption in a community are associated with greater measles incidence in that community, among both the exempt and nonexempt population

Pertussis

- In at least 7 statewide pertussis epidemics and multiple more geographically restricted outbreaks a substantial proportion of pertussis cases in certain age groups were unvaccinated or undervaccinated

- At least 7 pertussis outbreaks have been described in highly vaccinated populations, illustrative of waning vaccine-induced immunity
- Reasons for nonvaccination are infrequently reported in pertussis outbreaks, however in 8 outbreaks, between 59%–93% of the cases were intentionally unvaccinated or undervaccinated
- Individuals with vaccine exemptions have a greater risk for pertussis than those who are fully vaccinated
- Undervaccinated individuals—those who have received fewer than the recommended number of age-appropriate doses of pertussis-containing vaccine—also have an elevated risk for pertussis compared with fully vaccinated individuals
- Schools, communities, and states with higher exemption rates have higher rates of pertussis, including among the fully vaccinated population
- States with allowances for personal belief exemptions or with easier exemption policies have higher incidence rates of pertussis

Pertussis

Recent Epidemiology of Pertussis in the United States—Unlike measles, pertussis remains endemic in the United States. Following the widespread introduction of whole-cell pertussis vaccines into the routine childhood immunization schedule in the mid- 1940s, there was a marked reduction in pertussis incidence, from 150 000 to 260 000 cases annually to a nadir of 1010 cases in 1976.¹⁸ Since then, there has been a steady increase in the incidence of pertussis in the United States,^{51–53} with an interepidemic interval similar to the prevaccine era.⁵⁴ For the past 10 years, more than 10 000 cases have been reported annually (>25 000 in the last 3 years), matching figures that have not been observed in more than 5 decades (Box 1).¹⁸

This resurgence in pertussis has been attributed to multiple potential factors. Increased physician awareness and improvements in diagnostic techniques likely contributed to enhanced disease reporting early in the resurgence. Then, beginning in the mid-1990s, acellular pertussis vaccines replaced whole-cell vaccines in the childhood immunization schedule⁵⁵—although these acellular vaccines cause fewer local (eg, pain, swelling, and redness) and systemic (eg, fever and irritability) reactions,⁵⁶ they are less effective than whole-cell vaccines for preventing pertussis.⁵⁷ More recently, waning immunity to pertussis has been increasingly recognized as an important factor in disease resurgence.⁵⁸ Multiple observational studies, spanning epidemic and interepidemic periods, from a variety of geographic regions within the United States, found that a longer interval since receipt of the last pertussis-containing vaccine—including both the tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis (Tdap) booster for adolescents or DTaP—was associated with an increased risk of pertussis.^{59–64} A recent meta-analysis, which included studies within

and outside the United States, estimated that the odds of acquiring pertussis increased 1.33 times for each additional year since the last dose of childhood DTaP.⁶⁵

Vaccine Refusal and Pertussis Outbreaks—In the context of the now well-established association between waning immunity to pertussis and increased disease risk, we sought to identify the contribution of vaccine exemptions to outbreak-related pertussis. Because multiple doses of pertussis-containing vaccine are needed to develop and maintain protective immunity to pertussis, pertussis vaccination status can be classified as completely unvaccinated (ie, having received no doses of pertussis-containing vaccine) or undervaccinated (ie, having received fewer than the recommended number of age-appropriate doses of pertussis-containing vaccine). The Advisory Committee on Immunization Practices currently recommends 5 DTaP doses at age 2, 4, and 6 months, at 15 to 18 months, and again at 4 to 6 years, as well as a Tdap booster in adolescence (between age 11 and 12 years), or adulthood (if the adolescent dose was missed), and during every pregnancy.⁶⁶

We identified 1384 articles pertaining to pertussis outbreaks that have been described since the start of the pertussis resurgence (from January 1, 1977, through November 30, 2015). Of these, we screened 1123 unique titles and abstracts, and excluded 1028 from further review: these were studies conducted outside the United States, articles focusing on nonpertussis diseases, or papers deemed not relevant to the search query (eg, review articles, editorials, guideline or position statements, studies of vaccine coverage, immunogenicity, or safety, articles on general epidemiology that were not specific to pertussis, basic science studies, and studies on diagnosis, treatment, and prophylaxis of pertussis). After full-text review of the remaining 95 articles, 63 were excluded because they were duplicate reports of the same outbreak or they did not provide data relevant to this review, yielding 32 reports of nonoverlapping community or statewide pertussis outbreaks that we used for this summary (eFigure 2A in the Supplement).^{67–98} These reports included 10 609 pertussis cases among individuals for whom a vaccination status was reported (age range, 10 days–87 years).

In the 5 largest statewide epidemics, unvaccinated (or undervaccinated) individuals comprised a substantial proportion of cases in various age groups. For example, investigation of 781 cases during the Arizona pertussis epidemic of 1988 revealed that 33% of the 598 cases involving individuals who were age-eligible to have received at least 3 doses of pertussis-containing vaccine (ie, >6 months) were not appropriately immunized.⁷³ More recently, during the 2010 California pertussis epidemic, among the 4415 pertussis cases in children aged 6 months to 18 years for whom a vaccination history was available, 2001 children (45%) had not received age-appropriate pertussis-containing vaccines (380 unvaccinated children and 1621 undervaccinated children).⁸⁸ Similarly, in Washington in 2012, among 1829 cases in children aged 3 months to 19 years with available vaccination histories, 508 children (28%) were not up-to-date with their pertussis immunizations.⁸⁷ Oregon also experienced a pertussis epidemic in 2012, and among the pertussis cases aged 2 months to 6 years, 31% were unvaccinated and 24% were undervaccinated.⁹² Finally, during the most recent 2014 California pertussis epidemic, of the 222 cases among neonates and infants under the age of 12 months with detailed vaccine histories only 53 neonates and

infants (24%) had received any doses of DTaP, even though more than half (51%) of all the infants in this age group were age-eligible (older than 2 months) for pertussis vaccination.⁹³

Similar to the 5 statewide epidemics described above, there were 2 previous statewide outbreaks^{68,71} as well as multiple more geographically restricted outbreaks^{73,74,83,84,90,92,94} that also had substantial numbers of unvaccinated case patients; however, we also found reports of epidemics within populations with high rates of vaccination coverage,^{72,75,80,81,86,89,98} illustrating the important contribution of waning immunity to pertussis to disease epidemiology.^{86,98} Importantly, there was substantial heterogeneity among these reports in the criteria used to identify pertussis cases, the types of vaccine available for use at the time (ie, whole cell, acellular, and adult boosters), the proportion of cases in which an immunization history was available, the methods used to ascertain immunization histories, and how immunization status was described (eg, number of previous doses, above or below a threshold number of doses). In addition, owing to the inherent challenges of accurately identifying and diagnosing pertussis, chains of transmission could not be established in the vast majority of pertussis outbreaks. However, in 2 studies, secondary cases among fully vaccinated individuals, whose immunity had presumably waned and who likely acquired disease from intentionally unvaccinated individuals, were clearly identified.^{73,94}

Only 9 reports (describing 12 pertussis outbreaks) characterized the reason(s) that case patients were intentionally unvaccinated or undervaccinated—these reasons included personal belief (or philosophical) exemptions (70% of unvaccinated cases in Oregon in 2012⁹²), religious beliefs⁶⁸ or exemptions (84% of unvaccinated cases in Florida in 2013⁹⁴; 59%–93% of unvaccinated cases in 4 separate outbreaks in Massachusetts from 1986–1988⁷⁴), cultural norms (72% unvaccinated in an Amish community in Delaware in 2004–2005⁸⁴; 79% unvaccinated in an Amish community in Illinois in 2009–2010⁹⁰), illness at the time a vaccine dose was due,^{68,71} inability to appear for the vaccination appointment,⁷¹ or hesitance on the part of the vaccine provider.^{68,73}

Vaccine Refusal and Pertussis Risk—We identified 383 articles in the search for studies evaluating the association between nonmedical exemptions to routine vaccination and the individual or community risk for pertussis. Of these, 359 nonduplicate titles and abstracts were reviewed and 350 were excluded, yielding 9 studies that were included in this summary (eFigure 2B in the Supplement).^{10,12,50,99–104} The excluded papers included studies not specific to pertussis or to its epidemiology in the United States, review articles, opinion pieces, studies of vaccine effectiveness, immunogenicity, or safety, and articles on the basic science or clinical aspects of pertussis disease.

Three studies evaluated the individual risk of pertussis associated with vaccine refusal—1 retrospective cohort study used Colorado pertussis surveillance and immunization data from 1987–1998 and determined that those with exemptions were 5.9 times more likely to acquire pertussis compared with fully vaccinated individuals.⁵⁰ A different case-control study analyzed pertussis cases from 1996–2007 within a large managed care organization and computed a nearly 20-fold increased risk of pertussis among individuals with exemptions¹⁰⁰—11% of the pertussis cases in that cohort were attributed to vaccine refusal.

Another case-control study used pooled longitudinal data (2004–2010) from 8 Vaccine Safety Data link sites and determined that even undervaccinated individuals had an increased risk of pertussis, with the risk being proportional to the number of missed doses of DTaP.¹⁰²

We also found 6 observational studies that showed that high rates of vaccine exemption in a community or state are associated with an increased risk of pertussis in that community or state compared with communities or states without high rates of vaccine exemptions. For example, 4 different state-specific studies—in Colorado,⁵⁰ Michigan,⁹⁹ New York,¹⁰¹ and California (during the 2010 epidemic)¹²—all demonstrated that schools and communities with high vaccine exemption rates also had higher rates of pertussis. The risk for acquiring pertussis was higher even for those who were appropriately vaccinated. More generally, 3 different ecological studies found that states with allowances for personal belief exemptions¹⁰ or those with easier exemption processes^{10,104} have an elevated incidence of pertussis (approximately 1.5 times higher in 1 study¹⁰) compared with states with difficult exemption processes, an association that was confirmed using vaccine-specific uptake data.¹⁰³ This geospatial association between clusters of vaccine refusers and pertussis cases cannot be adequately explained by waning immunity because there should not be geographic heterogeneity in the duration of protection afforded by pertussis-containing vaccines. Together these data therefore suggest an association between high rates of vaccine exemption and the sustained transmission of vaccine-preventable diseases in the community (Box 2).

Discussion

In this review, unvaccinated or undervaccinated individuals comprised substantial proportions of cases in measles and some pertussis outbreaks, and vaccine refusal—as measured by population-level vaccine exemption rates—was associated with an elevated risk for measles and pertussis, including among fully vaccinated individuals. However, the available literature also suggests that vaccine refusal is not the only factor associated with outbreaks of vaccine-preventable diseases, and the relative importance of vaccine refusal varies by disease (eg, measles more than pertussis) and epidemiological context (eg, prevalence and density of individuals with exemptions in a community).

There are several limitations to our review. A formal assessment of study quality was not performed. Most, if not all, of the data on measles and pertussis outbreaks can be accessed using publicly available resources. We included several studies that analyzed measles and pertussis disease risk in the context of vaccine exemption rates, and such studies are subject to multiple important biases common to ecological, case-control, or cohort studies, including ascertainment and misclassification bias. However, these limitations would tend to bias estimates of disease risk toward the null (eg, classifying fully vaccinated individuals as exempted would only underestimate disease risk among those with exemptions). Thus the association between high rates of vaccine exemption and measles or pertussis risk is likely to be even greater than that reported in previous studies. In addition, given that the evidence for waning immunity to pertussis has been previously evaluated,⁶⁵ and its association with disease resurgence has been established,^{59–64} in this review we focused on the additional factor of vaccine refusal.

This review has broad implications for vaccine practice and policy. For instance, fundamental to the strength and legitimacy of justifications to override parental decisions to refuse a vaccine for their child is a clear demonstration that the risks and harms to the child of remaining unimmunized are substantial. Similarly, central to any justification to restrict individual freedom by mandating vaccines to prevent harm to others is an understanding of the nature and magnitude of these risks and harms. However, the risks of vaccine refusal remain imperfectly defined, and the association between vaccine refusal and vaccine-preventable diseases may be both population- and disease-specific. Vaccine refusal-specific strategies to optimize vaccine uptake could include state or school-level enforcement of vaccine mandates, or increasing the difficulty with which vaccine exemptions can be obtained. At the same time, immunization policy makers must also address the reasons for vaccine hesitancy, which may include parental perceptions regarding the risk and severity of vaccine-preventable diseases, the safety and effectiveness of routine immunizations, and confidence in medical professionals, corporations, and the health care system.¹⁰⁵ Recognizing the limitations of the existing evidence should not stifle practitioners' and policy makers' ability to counsel families and craft effective policy, but should serve as motivation to develop, refine, and improve disease surveillance, detection, and outcomes-based research.

Conclusions

A substantial proportion of the US measles cases in the era after elimination were intentionally unvaccinated. The phenomenon of vaccine refusal was associated with an increased risk for measles among people who refuse vaccines and among fully vaccinated individuals. Although pertussis resurgence has been attributed to waning immunity and other factors, vaccine refusal was still associated with an increased risk for pertussis in some populations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding/Support: This work is supported by award T32AI074492 from the Emory Vaccinology Training Program of the National Institute of Allergy and Infectious Diseases (Dr Phadke).

References

1. Elam-Evans LD, Yankey D, Singleton JA, Kolasa M. Centers for Disease Control and Prevention (CDC). National, state, and selected local area vaccination coverage among children aged 19–35 months—United States, 2013. *MMWR Morb Mortal Wkly Rep.* 2014; 63(34):741–748. [PubMed: 25166924]
2. Whitney CG, Zhou F, Singleton J, Schuchat A. Centers for Disease Control and Prevention (CDC). Benefits from immunization during the vaccines for children program era—United States, 1994–2013. *MMWR Morb Mortal Wkly Rep.* 2014; 63(16):352–355. [PubMed: 24759657]
3. Yang YT, Silverman RD. Legislative prescriptions for controlling nonmedical vaccine exemptions. *JAMA.* 2015; 313(3):247–248. [PubMed: 25602992]

4. Gostin LO. Law, ethics, and public health in the vaccination debates: politics of the measles outbreak. *JAMA*. 2015; 313(11):1099–1100. [PubMed: 25675396]
5. Yang YT, Barraza L, Weidenaar K. Measles outbreak as a catalyst for stricter vaccine exemption legislation. *JAMA*. 2015; 314(12):1229–1230. [PubMed: 26214397]
6. Centers for Disease Control and Prevention (CDC). Invasive *Haemophilus influenzae* type B disease in 5 young children—Minnesota, 2008. *MMWR Morb Mortal Wkly Rep*. 2009; 58(3):58–60. [PubMed: 19177041]
7. Glanz JM, McClure DL, Magid DJ, Daley MF, France EK, Hambidge SJ. Parental refusal of varicella vaccination and the associated risk of varicella infection in children. *Arch Pediatr Adolesc Med*. 2010; 164(1):66–70. [PubMed: 20048244]
8. Glanz JM, McClure DL, O’Leary ST, et al. Parental decline of pneumococcal vaccination and risk of pneumococcal related disease in children. *Vaccine*. 2011; 29(5):994–999. [PubMed: 21145372]
9. Salmon DA, Haber M, Gangarosa EJ, Phillips L, Smith NJ, Chen RT. Health consequences of religious and philosophical exemptions from immunization laws: individual and societal risk of measles. *JAMA*. 1999; 282(1):47–53. [PubMed: 10404911]
10. Omer SB, Pan WK, Halsey NA, et al. Nonmedical exemptions to school immunization requirements: secular trends and association of state policies with pertussis incidence. *JAMA*. 2006; 296(14):1757–1763. [PubMed: 17032989]
11. Omer SB, Richards JL, Ward M, Bednarczyk RA. Vaccination policies and rates of exemption from immunization, 2005–2011. *N Engl J Med*. 2012; 367(12):1170–1171. [PubMed: 22992099]
12. Atwell JE, Van Otterloo J, Zipprich J, et al. Nonmedical vaccine exemptions and pertussis in California, 2010. *Pediatrics*. 2013; 132(4):624–630. [PubMed: 24082000]
13. Majumder MS, Cohn EL, Mekaru SR, Huston JE, Brownstein JS. Substandard vaccination compliance and the 2015 measles outbreak. *JAMA Pediatr*. 2015; 169(5):494–495. [PubMed: 25774618]
14. Clemmons NS, Gastanaduy PA, Fiebelkorn AP, Redd SB, Wallace GS. Centers for Disease Control and Prevention (CDC). Measles—United States, January 4–April 2, 2015. *MMWR Morb Mortal Wkly Rep*. 2015; 64(14):373–376. [PubMed: 25879894]
15. Centers for Disease Control and Prevention. [Accessed July 17, 2015] Measles cases and outbreaks. <http://www.cdc.gov/measles/cases-outbreaks.html>
16. Zipprich J, Winter K, Hacker J, Xia D, Watt J, Harriman K. Centers for Disease Control and Prevention (CDC). Measles outbreak—California, December 2014–February 2015. *MMWR Morb Mortal Wkly Rep*. 2015; 64(6):153–154. [PubMed: 25695321]
17. Katz SL, Hinman AR. Summary and conclusions: measles elimination meeting, 16–17 March 2000. *J Infect Dis*. 2004; 189(suppl 1):S43–S47. [PubMed: 15106088]
18. [Accessed March 25, 2015] Surveillance & Reporting. Pertussis (whooping cough). <http://www.cdc.gov/pertussis/surv-reporting.html>
19. Hamborsky, J.; Kroger, A.; Wolfe, S., editors. Measles: Epidemiology and Prevention of Vaccine-Preventable Diseases (The Pink Book). 13. Washington, DC: Public Health Foundation; 2015.
20. Centers for Disease Control (CDC). Public-sector vaccination efforts in response to the resurgence of measles among preschool-aged children—United States, 1989–1991. *MMWR Morb Mortal Wkly Rep*. 1992; 41(29):522–525. [PubMed: 1630430]
21. Centers for Disease Control and Prevention (CDC). Measles—United States, 2000. *MMWR Morb Mortal Wkly Rep*. 2002; 51(6):120–123. [PubMed: 11898926]
22. Centers for Disease Control and Prevention (CDC). Epidemiology of measles—United States, 2001–2003. *MMWR Morb Mortal Wkly Rep*. 2004; 53(31):713–716. [PubMed: 15306756]
23. Centers for Disease Control and Prevention (CDC). Measles—United States, 2004. *MMWR Morb Mortal Wkly Rep*. 2005; 54(48):1229–1231. [PubMed: 16340938]
24. Centers for Disease Control and Prevention (CDC). Measles—United States, 2005. *MMWR Morb Mortal Wkly Rep*. 2006; 55(50):1348–1351. [PubMed: 17183226]
25. Centers for Disease Control and Prevention (CDC). Measles among adults associated with adoption of children in China—California, Missouri, and Washington, July–August 2006. *MMWR Morb Mortal Wkly Rep*. 2007; 56(7):144–146. [PubMed: 17318114]

26. Centers for Disease Control and Prevention (CDC). Multistate measles outbreak associated with an international youth sporting event— Pennsylvania, Michigan, and Texas, August–September 2007. *MMWR Morb Mortal Wkly Rep.* 2008; 57(7):169–173. [PubMed: 18288074]
27. Centers for Disease Control and Prevention (CDC). Update: measles—United States, January–July 2008. *MMWR Morb Mortal Wkly Rep.* 2008; 57(33):893–896. [PubMed: 18716580]
28. Chen SY, Anderson S, Kutty PK, et al. Health care-associated measles outbreak in the United States after an importation: challenges and economic impact. *J Infect Dis.* 2011; 203(11):1517–1525. [PubMed: 21531693]
29. Centers for Disease Control and Prevention (CDC). Hospital-associated measles outbreak— Pennsylvania, March–April 2009. *MMWR Morb Mortal Wkly Rep.* 2012; 61(2):30–32. [PubMed: 22258416]
30. Coleman MS, Garbat-Welch L, Burke H, et al. Direct costs of a single case of refugee-imported measles in Kentucky. *Vaccine.* 2012; 30(2):317–321. [PubMed: 22085555]
31. Centers for Disease Control and Prevention (CDC). Measles among US-bound refugees from Malaysia—California, Maryland, North Carolina, and Wisconsin, August–September 2011. *MMWR Morb Mortal Wkly Rep.* 2011; 60(37):1281–1282. [PubMed: 21937975]
32. Centers for Disease Control and Prevention (CDC). Measles outbreak associated with an arriving refugee—Los Angeles County, California, August–September 2011. *MMWR Morb Mortal Wkly Rep.* 2012; 61(21):385–389. [PubMed: 22647743]
33. Centers for Disease Control and Prevention (CDC). Measles—United States, 2011. *MMWR Morb Mortal Wkly Rep.* 2012; 61:253–257. [PubMed: 22513526]
34. Centers for Disease Control and Prevention (CDC). Measles—United States, January 1–August 24, 2013. *MMWR Morb Mortal Wkly Rep.* 2013; 62(36):741–743. [PubMed: 24025755]
35. Nyangoma EN, Olson CK, Benoit SR, et al. Centers for Disease Control and Prevention (CDC). Measles outbreak associated with adopted children from China—Missouri, Minnesota, and Washington, July 2013. *MMWR Morb Mortal Wkly Rep.* 2014; 63(14):301–304. [PubMed: 24717816]
36. Gastañaduy PA, Redd SB, Fiebelkorn AP, et al. Division of Viral Disease, National Center for Immunization and Respiratory Diseases, CDC. Measles—United States, January 1–May 23, 2014. *MMWR Morb Mortal Wkly Rep.* 2014; 63(22):496–499. [PubMed: 24898167]
37. Slade TA, Klekamp B, Rico E, Mejia-Echeverry A. Measles outbreak in an unvaccinated family and a possibly associated international traveler—Orange County, Florida, December 2012–January 2013. *MMWR Morb Mortal Wkly Rep.* 2014; 63(36):781–784. [PubMed: 25211542]
38. Centers for Disease Control and Prevention (CDC). Outbreak of measles—San Diego, California, January–February 2008. *MMWR Morb Mortal Wkly Rep.* 2008; 57(8):203–206. [PubMed: 18305451]
39. Bandyopadhyay AS, Bandy U. Emerging global epidemiology of measles and public health response to confirmed case in Rhode Island. *R I Med J* (2013). 2013; 96(2):41–44. [PubMed: 23641427]
40. Centers for Disease Control and Prevention (CDC). Measles outbreak in a boarding school— Pennsylvania, 2003. *MMWR Morb Mortal Wkly Rep.* 2004; 53(14):306–309. [PubMed: 15085075]
41. Centers for Disease Control and Prevention (CDC). Measles outbreak associated with an imported case in an infant—Alabama, 2002. *MMWR Morb Mortal Wkly Rep.* 2004; 53(2):30–33. [PubMed: 14737061]
42. Centers for Disease Control and Prevention (CDC). Post exposure prophylaxis, isolation, and quarantine to control an import-associated measles outbreak—Iowa, 2004. *MMWR Morb Mortal Wkly Rep.* 2004; 53(41):969–971. [PubMed: 15496826]
43. Centers for Disease Control and Prevention (CDC). Update: Multistate investigation of measles among adoptees from China—April 16, 2004. *MMWR Morb Mortal Wkly Rep.* 2004; 53(15): 323–324. [PubMed: 15103298]
44. Centers for Disease Control and Prevention (CDC). Two measles outbreaks after importation— Utah, March–June 2011. *MMWR Morb Mortal Wkly Rep.* 2013; 62(12):222–225. [PubMed: 23535688]

45. Gahr P, DeVries AS, Wallace G, et al. An outbreak of measles in an undervaccinated community. *Pediatrics*. 2014; 134(1):e220–e228. [PubMed: 24913790]
46. Rosen JB, Rota JS, Hickman CJ, et al. Outbreak of measles among persons with prior evidence of immunity, New York City, 2011. *Clin Infect Dis*. 2014; 58(9):1205–1210. [PubMed: 24585562]
47. Vega JS, Escobedo M, Schulte CR, et al. Centers for Disease Control and Prevention (CDC). Notes from the field: measles transmission at a domestic terminal gate in an international airport—United States, January 2014. *MMWR Morb Mortal Wkly Rep*. 2014; 63(50):1211. [PubMed: 25522093]
48. Banerjee E, Hickman C, Engels K, Kenyon C. Centers for Disease Control and Prevention (CDC). Notes from the field: measles transmission in an international airport at a domestic terminal gate—April–May 2014. *MMWR Morb Mortal Wkly Rep*. 2015; 64(24):679. [PubMed: 26110839]
49. Collier MG, Cierzniewski A, Duszynski T, et al. Measles outbreak associated with international travel, Indiana, 2011. *J Pediatric Infect Dis Soc*. 2013; 2(2):110–118. [PubMed: 26619458]
50. Feikin DR, Lezotte DC, Hamman RF, Salmon DA, Chen RT, Hoffman RE. Individual and community risks of measles and pertussis associated with personal exemptions to immunization. *JAMA*. 2000; 284(24):3145–3150. [PubMed: 11135778]
51. Farizo KM, Cochi SL, Zell ER, Brink EW, Wassilak SG, Patriarca PA. Epidemiological features of pertussis in the United States, 1980–1989. *Clin Infect Dis*. 1992; 14(3):708–719. [PubMed: 1562663]
52. Güri D, Strebel PM, Bardenheier B, et al. Changing epidemiology of pertussis in the United States: increasing reported incidence among adolescents and adults, 1990–1996. *Clin Infect Dis*. 1999; 28(6):1230–1237. [PubMed: 10451158]
53. Centers for Disease Control and Prevention (CDC). Resurgence of pertussis—United States, 1993. *MMWR Morb Mortal Wkly Rep*. 1993; 42(49):952–953. 959–960. [PubMed: 8246860]
54. Cherry JD. The epidemiology of pertussis and pertussis immunization in the United Kingdom and the United States: a comparative study. *Curr Probl Pediatr*. 1984; 14(2):1–78. [PubMed: 6368134]
55. Pertussis vaccination: use of acellular pertussis vaccines among infants and young children: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Morb Mortal Wkly Rep*. 1997; 46(RR-7):1–25. [PubMed: 9011775]
56. Decker MD, Edwards KM, Steinhoff MC, et al. Comparison of 13 acellular pertussis vaccines: adverse reactions. *Pediatrics*. 1995; 96(3 pt 2):557–566. [PubMed: 7659476]
57. Edwards, KM.; Decker, MD. Pertussis vaccines. In: Orenstein, WA.; Offit, PA., editors. *Vaccines*. 6. Plotkin, SA: Elsevier Saunders; 2013.
58. Versteegh FG, Schellekens JF, Nagelkerke AF, Roord JJ. Laboratory-confirmed reinfections with *Bordetella pertussis*. *Acta Paediatr*. 2002; 91(1):95–97. [PubMed: 11885549]
59. Lavine J, Broutin H, Harvill ET, Bjørnstad ON. Imperfect vaccine-induced immunity and whooping cough transmission to infants. *Vaccine*. 2010; 29(1):11–16. [PubMed: 21034823]
60. Witt MA, Katz PH, Witt DJ. Unexpectedly limited durability of immunity following acellular pertussis vaccination in preadolescents in a North American outbreak. *Clin Infect Dis*. 2012; 54(12):1730–1735. [PubMed: 22423127]
61. Klein NP, Bartlett J, Rowhani-Rahbar A, Fireman B, Baxter R. Waning protection after fifth dose of acellular pertussis vaccine in children. *N Engl J Med*. 2012; 367(11):1012–1019. [PubMed: 22970945]
62. Tartof SY, Lewis M, Kenyon C, et al. Waning immunity to pertussis following 5 doses of DTaP. *Pediatrics*. 2013; 131(4):e1047–e1052. [PubMed: 23478868]
63. Misegades LK, Winter K, Harriman K, et al. Association of childhood pertussis with receipt of 5 doses of pertussis vaccine by time since last vaccine dose, California, 2010. *JAMA*. 2012; 308(20):2126–2132. [PubMed: 23188029]
64. Koepke R, Eickhoff JC, Ayele RA, et al. Estimating the effectiveness of tetanus, diphtheria, acellular pertussis vaccine (Tdap) for preventing pertussis: evidence of rapidly waning immunity and difference in effectiveness by Tdap brand. *J Infect Dis*. 2014; 210(6):942–953. [PubMed: 24903664]
65. McGirr A, Fisman DN. Duration of pertussis immunity after DTaP immunization: a meta-analysis. *Pediatrics*. 2015; 135(2):331–343. [PubMed: 25560446]

66. Advisory Committee on Immunization Practices. [Accessed October 12, 2015] Recommended immunization schedules for persons aged 0 through 18 years. <http://www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-child-combined-schedule.pdf>
67. Broome CV, Preblud SR, Bruner B, et al. Epidemiology of pertussis, Atlanta, 1977. *J Pediatr*. 1981; 98(3):362–367. [PubMed: 7009816]
68. Centers for Disease Control (CDC). Pertussis—Maryland, 1982. *MMWR Morb Mortal Wkly Rep*. 1983; 32(23):297–300. 305. [PubMed: 6406816]
69. Leads from the MMWR. Leads from the MMWR Pertussis—Washington, 1984. *JAMA*. 1985; 254(5):600–602. [PubMed: 3892068]
70. Vichyanond P, Olson LC. Pertussis outbreak in SE Kansas—1983–1984. *Kans Med*. 1985; 86(2): 52–54. 62. [PubMed: 3981853]
71. Nkowane BM, Wassilak SG, McKee PA, et al. Pertussis epidemic in Oklahoma: difficulties in preventing transmission. *Am J Dis Child*. 1986; 140(5):433–437. [PubMed: 3962936]
72. Biellik RJ, Patriarca PA, Mullen JR, et al. Risk factors for community- and household-acquired pertussis during a large-scale outbreak in central Wisconsin. *J Infect Dis*. 1988; 157(6):1134–1141. [PubMed: 3373018]
73. Ey JL, Duncan B, Barton LL, Buckett G. The influence of preschool pertussis immunization on an epidemic of pertussis. *Pediatr Infect Dis J*. 1991; 10(8):576–578. [PubMed: 1891289]
74. Etkind P, Lett SM, Macdonald PD, Silva E, Peppe J. Pertussis outbreaks in groups claiming religious exemptions to vaccinations. *Am J Dis Child*. 1992; 146(2):173–176. [PubMed: 1733146]
75. Centers for Disease Control and Prevention (CDC). Pertussis outbreaks—Massachusetts and Maryland, 1992. *MMWR Morb Mortal Wkly Rep*. 1993; 42(11):197–200. [PubMed: 8446095]
76. Christie CD, Marx ML, Marchant CD, Reising SF. The 1993 epidemic of pertussis in Cincinnati: resurgence of disease in a highly immunized population of children. *N Engl J Med*. 1994; 331(1): 16–21. [PubMed: 8202096]
77. Davis SF, Sutter RW, Strebel PM, et al. Concurrent outbreaks of pertussis and *Mycoplasma pneumoniae* infection: clinical and epidemiological characteristics of illnesses manifested by cough. *Clin Infect Dis*. 1995; 20(3):621–628. [PubMed: 7756486]
78. Kenyon TA, Izurieta H, Shulman ST, et al. Large outbreak of pertussis among young children in Chicago, 1993: investigation of potential contributing factors and estimation of vaccine effectiveness. *Pediatr Infect Dis J*. 1996; 15(8):655–661. [PubMed: 8858667]
79. Centers for Disease Control and Prevention (CDC). Pertussis outbreak—Vermont, 1996. *MMWR Morb Mortal Wkly Rep*. 1997; 46(35):822–826. [PubMed: 9310216]
80. Brennan M, Strebel P, George H, et al. Evidence for transmission of pertussis in schools, Massachusetts, 1996: epidemiologic data supported by pulsed-field gel electrophoresis studies. *J Infect Dis*. 2000; 181(1):210–215. [PubMed: 10608769]
81. Khetsuriani N, Bisgard K, Prevots DR, et al. Pertussis outbreak in an elementary school with high vaccination coverage. *Pediatr Infect Dis J*. 2001; 20(12):1108–1112. [PubMed: 11740314]
82. Centers for Disease Control and Prevention (CDC). School-associated pertussis outbreak—Yavapai County, Arizona, September 2002–February 2003. *MMWR Morb Mortal Wkly Rep*. 2004; 53(10): 216–219. [PubMed: 15029116]
83. Khan FN, Lin M, Hinkle CJ, et al. Case-control study of vaccination history in relation to pertussis risk during an outbreak among school students. *Pediatr Infect Dis J*. 2006; 25(12):1132–1136. [PubMed: 17133158]
84. Centers for Disease Control and Prevention (CDC). Pertussis outbreak in an Amish community—Kent County, Delaware, September 2004–February 2005. *MMWR Morb Mortal Wkly Rep*. 2006; 55(30):817–821. [PubMed: 16888610]
85. Schafer S, Gillette H, Hedberg K, Cieslak P. A community-wide pertussis outbreak: an argument for universal booster vaccination. *Arch Intern Med*. 2006; 166(12):1317–1321. [PubMed: 16801516]
86. Sotir MJ, Cappozzo DL, Warshauer DM, et al. A countywide outbreak of pertussis: initial transmission in a high school weight room with subsequent substantial impact on adolescents and adults. *Arch Pediatr Adolesc Med*. 2008; 162(1):79–85. [PubMed: 18180417]

87. Centers for Disease Control and Prevention (CDC). Pertussis epidemic—Washington, 2012. *MMWR Morb Mortal Wkly Rep.* 2012; 61(28):517–522. [PubMed: 22810264]
88. Winter K, Harriman K, Zipprich J, et al. California pertussis epidemic, 2010. *J Pediatr.* 2012; 161(6):1091–1096. [PubMed: 22819634]
89. Bass JB Jr, Turpin-Saunders SR. Coughing up answers: a community's response to pertussis. *NC Med J.* 2013; 74(5):420–424.
90. Medina-Marino A, Reynolds D, Finley C, Hays S, Jones J, Soyemi K. Communication and mass vaccination strategies after pertussis outbreak in rural Amish communities-Illinois, 2009–2010. *J Rural Health.* 2013; 29(4):413–419. [PubMed: 24088215]
91. Rodgers L, Martin SW, Cohn A, et al. Epidemiologic and laboratory features of a large outbreak of pertussis-like illnesses associated with cocirculating *Bordetella holmesii* and *Bordetella pertussis*—Ohio, 2010–2011. *Clin Infect Dis.* 2013; 56(3):322–331. [PubMed: 23087388]
92. Liko J, Robison SG, Cieslak PR. Pertussis vaccine performance in an epidemic year—Oregon, 2012. *Clin Infect Dis.* 2014; 59(2):261–263. [PubMed: 24748523]
93. Winter K, Glaser C, Watt J, Harriman K. Centers for Disease Control and Prevention (CDC). Pertussis epidemic—California, 2014. *MMWR Morb Mortal Wkly Rep.* 2014; 63(48):1129–1132. [PubMed: 25474033]
94. Matthias J, Dusek C, Pritchard SP, Rutledge L, Kinchen P, Lander M. Centers for Disease Control and Prevention. Notes from the field: outbreak of pertussis in a school and religious community averse to health care and vaccinations—Columbia County, Florida, 2013. *MMWR Morb Mortal Wkly Rep.* 2014; 63(30):655. [PubMed: 25078655]
95. Theofiles AG, Cunningham SA, Chia N, et al. Pertussis outbreak, southeastern Minnesota, 2012. *Mayo Clin Proc.* 2014; 89(10):1378–1388. [PubMed: 25282430]
96. Gao F, Mahoney JC, Daly ER, Lamothe W, Tullo D, Bean C. Evaluation of a multitarget real-time PCR assay for detection of *Bordetella* species during a pertussis outbreak in New Hampshire in 2011. *J Clin Microbiol.* 2014; 52(1):302–306. [PubMed: 24131698]
97. Cantey JB, Sanchez PJ, Tran J, Chung W, Siegel JD. Pertussis: a persistent cause of morbidity and mortality in young infants. *J Pediatr.* 2014; 164(6):1489–1492. e1. [PubMed: 24565424]
98. Eshofonie AO, Lin H, Valcin RP, Martin LR, Grunenwald PE. An outbreak of pertussis in rural Texas: an example of the resurgence of the disease in the United States. *J Community Health.* 2015; 40(1):88–91. [PubMed: 24927977]
99. Omer SB, Enger KS, Moulton LH, Halsey NA, Stokley S, Salmon DA. Geographic clustering of nonmedical exemptions to school immunization requirements and associations with geographic clustering of pertussis. *Am J Epidemiol.* 2008; 168(12):1389–1396. [PubMed: 18922998]
100. Glanz JM, McClure DL, Magid DJ, et al. Parental refusal of pertussis vaccination is associated with an increased risk of pertussis infection in children. *Pediatrics.* 2009; 123(6):1446–1451. [PubMed: 19482753]
101. Imdad A, Tserenpuntsag B, Blog DS, Halsey NA, Easton DE, Shaw J. Religious exemptions for immunization and risk of pertussis in New York State, 2000–2011. *Pediatrics.* 2013; 132(1):37–43. [PubMed: 23733795]
102. Glanz JM, Narwaney KJ, Newcomer SR, et al. Association between undervaccination with diphtheria, tetanus toxoids, and acellular pertussis (DTaP) vaccine and risk of pertussis infection in children 3 to 36 months of age. *JAMA Pediatr.* 2013; 167(11):1060–1064. [PubMed: 24019039]
103. Yang YT, Debold V. A longitudinal analysis of the effect of nonmedical exemption law and vaccine uptake on vaccine-targeted disease rates. *Am J Public Health.* 2014; 104(2):371–377. [PubMed: 24328666]
104. Bradford WD, Mandich A. Some state vaccination laws contribute to greater exemption rates and disease outbreaks in the United States. *Health Aff (Millwood).* 2015; 34(8):1383–1390. [PubMed: 26240253]
105. Salmon DA, Dudley MZ, Glanz JM, Omer SB. Vaccine hesitancy: causes, consequences, and a call to action. *Am J Prev Med.* 2015; 49(6 suppl 4):S391–S398. [PubMed: 26337116]




Figure 1. Cumulative Epidemic Curve of 18 Measles Outbreaks From 2000 Through 2015^a

Day 0 indicates index cases. Median cases per outbreak, 6.5 (range, 1–22).

^aThe cumulative epidemic curve presents data of measles outbreaks for which individual-level day of symptom onset and vaccination status was available (N = 145).

Characteristics of Measles Outbreak and Summary Reports Used to Evaluate Vaccine Refusals Among Measles Cases in the Era After Elimination

Table 1

Type of Report	Location	Year(s)	Measles Cases, No.	Vaccination Status Reported	Vaccine Status of Individuals, No.			
					Vaccinated	Unvaccinated	Unknown	
Summary ²¹	Nationwide	2000	86	86	23	40	NA	23
Summary ²²	Nationwide	2001–2003	216	216	42	174 ^b	NA	0
Summary ²³	Nationwide	2004	37	35	4	26	NA	7
Summary ²⁴	Nationwide	2005	66	66	8	50	NA	8
Outbreak ²⁵	Missouri, California, Washington	2006	3	3	3	0	NA	0
Outbreak ²⁶	Pennsylvania, Michigan, Texas	2007	7	7	2	3	NA	2
Summary ²⁷	Nationwide	2008	131	123	11	112	63	0
Outbreak ²⁸	Arizona	2008	15	15	0	8	NA	6
Outbreak ²⁹	Pennsylvania	2009	6	6	2	3	2	1
Outbreak ³⁰	Kentucky	2010	1	1	0	0	NA	1
Outbreak ³¹	California, Maryland, North Carolina, Wisconsin	2011	8	8	0	6	NA	2
Outbreak ³²	California	2011	4	4	1	3	NA	0
Summary ³³	Nationwide	2011	222	222	31	144	50	47
Summary ³⁴	16 US states	2013	159	159	13	131	92	15
Outbreak ³⁵	Missouri, Minnesota, Washington	2013	4	4	1	3	NA	0
Summary ³⁶	Nationwide	2014	288	288	30	200	165	58
Outbreak ³⁷	Florida	2014	5	5	0	4	4	1
Outbreak ¹⁴	Nationwide	2014–2015	159	99	28	71	29	60
Total	United States	2000–2015	1416	1408	199	804	405	231

Abbreviations: NA, not available; NME, nonmedical exemption.

^aThis column is a subset of the unvaccinated column.

^bThe 2001–2003 report²² did not distinguish between unvaccinated and unknown vaccination status, therefore the total at the bottom of the column does not include these 174 individuals.

Table 2

Characteristics of Studies Used to Evaluate Measles Risk Associated With Vaccine Refusal

Source	Study Design	Period	Location	Data Sources
Salmon et al, ⁹ 1999	Population-based retrospective cohort study to determine the individual risk for measles among exempt and fully vaccinated individuals; mathematical modeling study to determine the community risk of measles associated with clustering of exempt individuals	1985–1992	Nationwide	CDC Measles Surveillance System and annual state immunization program reports
Feikin et al, ⁵⁰ 2000	Population-based retrospective cohort study to determine the individual risk for measles (and pertussis) among exempt and fully vaccinated individuals, as well as the association between county-level measles incidence and school-based outbreaks with the rate of vaccine exemptions	1987–1998	Colorado	Colorado Department of Public Health and Environment and annual state immunization program reports

Abbreviation: CDC, Centers for Disease Control and Prevention.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript